



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

fever, dysentery, whooping cough, cholera morbus, smallpox, and measles were less prevalent than in the preceding week. Meningitis was reported present at 1, whooping cough at 8, measles at 14, smallpox at 24, diphtheria at 44, scarlet fever at 59, enteric fever at 80, and phthisis pulmonalis at 184 places.

The Monthly Bulletin says: There were 2,624 deaths reported to the Department of State for the month of October, or 98 less than the number returned for the preceding month. The death rate was 12.5 per 1,000 population. Deaths by ages were as follows: Under 1 year, 489; 1 to 4 years, inclusive, 185; 65 years and over, 766. There was a considerable decrease in deaths of infants and children, and a slight increase in deaths of elderly persons, as compared with September.

Important causes of death were as follows: Tuberculosis of the lungs, 190; other forms of tuberculosis, 35; typhoid fever, 90; diphtheria and croup, 78; scarlet fever, 7; measles, 3; whooping cough, 16; pneumonia, 143; diarrhea and enteritis, under two years, 173; cancer, 129; accidents and violence, 202.

There was 1 death from smallpox, which occurred in the city of Hillsdale. Somewhat increased mortality was shown from tuberculosis, diphtheria and croup, and pneumonia. Typhoid fever caused 90 deaths, as compared with 74 for the preceding month. There was a very large decrease in deaths reported from diarrheal diseases of children, and also a considerable diminution in the number of deaths from cancer.

Both urban and rural populations show less than the average mortality from typhoid fever for the month of October. Cities with unusually high death rates were Grand Rapids, Bay City, Flint, Kalamazoo, Lansing, Port Huron, West Bay City, referring only to cities of 10,000 population or upwards.

MINNESOTA—*Duluth*.—Month of October, 1903. Census population, 52,969. Total number of deaths, 55, including diphtheria 1, enteric fever 9, and 6 from tuberculosis.

UTAH—*Salt Lake City*.—Month of October, 1903. Estimated population, 75,000. Total number of deaths, 50, including enteric fever 6, and 1 from tuberculosis.

### *Report of immigration at Baltimore.*

OFFICE OF THE COMMISSIONER OF IMMIGRATION,  
*Baltimore, Md., November 28, 1903.*

*Number of alien immigrants who arrived at this port during the week ended November 28, 1903, also name of vessel and port from which they came.*

Date of arrival.	Name of vessel.	Where from.	Number of immigrants.
Nov. 26	Frankfurt .....	Bremen .....	1,171
	Total .....	.....	1,171

LOUIS T. WEIS, *Commissioner.*